

URINARY U/C



Available in 2 kg, 7.5 kg, 14 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
	2	55	5/8	48	4/8	42	4/8
	4	92	1	81	7/8	70	6/8
	6	125	1 + 3/8	110	1 + 2/8	95	1
	8	155	1 + 6/8	137	1 + 4/8	118	1 + 2/8
	10	184	2	162	1 + 6/8	140	1 + 4/8
	15	249	2 + 6/8	219	2 + 3/8	189	2 + 1/8
	20	309	3 + 3/8	272	3	235	2 + 5/8
	25	365	4	321	3 + 4/8	278	3
	30	419	4 + 5/8	368	4	318	3 + 4/8
	35	470	5 + 1/8	414	4 + 4/8	357	3 + 7/8
	40	520	5 + 6/8	457	5	395	4 + 3/8
	45	568	6 + 2/8	499	5 + 4/8	431	4 + 6/8
	50	614	6 + 6/8	541	5 + 7/8	467	5 + 1/8
	55	660	7 + 2/8	581	6 + 3/8	501	5 + 4/8
	60	704	7 + 6/8	620	6 + 6/8	535	5 + 7/8
	70	791	8 + 5/8	696	7 + 5/8	601	6 + 5/8
	80	874	9 + 5/8	769	8 + 3/8	664	7 + 2/8

Urinary U/C is a complete dietetic feed for dogs, formulated to reduce the formation of urate and cystine stones through its low level of protein and of purines, its high quality protein, its moderate level of sulphur amino acids and its urine alkalisating properties.

RECOMMENDATIONS: Feed Urinary U/C for up to 6 months to reduce the formation of urate stones, for up to 1 year to reduce the formation of cystine stones and for life in the case of irreversible disturbance of uric acid metabolism. It is recommended that a veterinarian's opinion be sought before use or before extending the period of use. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

LOWER URINARY TRACT DISEASE



RECOMMENDED FOR CASES OF

- Metabolic uroliths (urate, xanthine, cystine): management of recurrence
- Hyperuricosuria
- Management of metabolic uroliths risk in dogs with leishmaniasis treated with xanthine oxidase inhibitors eg allopurinol (if they present a secondary kidney disease (protein losing nephropathy with or without azotemia, then switch to a Renal diet)

NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



URATE CONTROL

A selection of proteins with low purine content helps reduce the formation of urate urinary stones.



CYSTINE CONTROL

A limited intake of proteins and of some amino acids helps limit the formation of cystine urinary stones.



SKIN BARRIER

A complex to support the barrier effect of the skin.



ANTIOXIDANT COMPLEX

A synergistic complex of antioxidants to help neutralise free radicals.

KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	47.03 g	18 g
Fat content	39.2 g	15 g
Carbohydrate	130.65 g	50 g
Crude fibre	5.75 g	2.2 g
Dietary fibre	14.89 g	5.7 g
Calcium	2.06 g	0.79 g
Phosphorus	1.54 g	0.59 g
Potassium	2.35 g	0.9 g
Sodium	0.78 g	0.3 g
Chloride	1.44 g	0.55 g
Magnesium	0.1 g	0.04 g
Copper	3.92 mg	1.5 mg
Zinc	44.42 mg	17 mg
Taurine	0.55 g	0.21 g
Vitamin A	7839.04 IU	3000 IU
Vitamin E	159.39 mg	61 mg
Omega 6	7.24 g	2.77 g
Omega 3	1.59 g	0.61 g
EPA + DHA	0.68 g	0.26 g
Metabolisable energy*	383	

*Calculated according to NRC2006.

COMPOSITION

Rice, maize flour, wheat flour, animal fats, egg powder, wheat gluten**, maize gluten, minerals, vegetable fibres, fish oil, soya oil, fructo-oligo-saccharides, borage oil, marigold extract (source of lutein). Protein sources: egg powder, wheat gluten, maize gluten.

**L.I.P.: protein selected for its very high digestibility.

ADDITIVES (KG)***

Nutritional additives: Vitamin A: 30000 IU, Vitamin D3: 800 IU, E1 (Iron): 63 mg, E2 (Iodine): 5.3 mg, E4 (Copper): 14 mg, E5 (Manganese): 66 mg, E6 (Zinc): 147 mg, E8 (Selenium): 0.16 mg - Preservatives - Antioxidants.

***Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY U/C

