

# URINARY S/O



THIN SLICES IN GRAVY

Available in 100 g

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

DOG'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	pouch	grams	pouch	grams	pouch
<b>2</b>	207	2	182	2	157	1 +1/2
<b>4</b>	348	3+1/2	306	3	264	2 +1/2
<b>6</b>	471	4 +1/2	415	4	358	3 +1/2
<b>8</b>	585	6	515	5	444	4 +1/2
<b>10</b>	691	7	608	6	525	5 +1/2
<b>15</b>	937	9 +1/2	824	8	712	7
<b>20</b>	1162	11 +1/2	1023	10	883	9
<b>25</b>	1374	13 +1/2	1209	12	1044	10 +1/2
<b>30</b>	1576	16	1386	14	1197	12
<b>35</b>	1769	17 +1/2	1556	15 +1/2	1344	13 +1/2
<b>40</b>	1955	19 +1/2	1720	17	1486	15

Urinary S/O is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

**IN OLDER DOGS:** the use of Urinary S/O Ageing\* is recommended and renal function should be checked before the advice of a new diet. It is recommended to divide the daily intake into 2 meals.

## LOWER URINARY TRACT DISEASE



### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



#### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



#### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.

**[S/O]® INDEX** This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	86.11 g	8 g
Fat content	59.2 g	5.5 g
Carbohydrate	51.67 g	4.8 g
Crude fibre	12.92 g	1.2 g
Dietary fibre	15.07 g	1.4 g
Calcium	1.61 g	0.15 g
Phosphorus	1.51 g	0.14 g
Potassium	1.61 g	0.15 g
Sodium	2.91 g	0.27 g
Chloride	3.98 g	0.37 g
Magnesium	0.13 g	0.01 g
Copper	3.77 mg	0.35 mg
Zinc	23.14 mg	2.15 mg
Taurine	1.24 g	0.12 g
Vitamin A	46286.33 IU	4300 IU
Vitamin E	226.05 mg	21 mg
Omega 6	19.38 g	1.8 g
Omega 3	1.35 g	0.13 g
EPA + DHA	0,65	0.06 g
Metabolisable energy*		93

\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, cereals, oils and fats, vegetable protein extracts, derivatives of vegetable origin, minerals, various sugars.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 210 IU, E1 (Iron): 3.5 mg, E2 (Iodine): 0.25 mg, E4 (Copper): 2 mg, E5 (Manganese): 1.1 mg, E6 (Zinc): 11 mg, Sulfur: 0.2%, Urine acidifying substances: calcium sulfate: 0.25%.  
Technological additives: (acidity regulator) Sodium bisulfate: 1.107 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

